

AN EXACTLY SOLVABLE CONSTRAINED XXZ CHAIN*

F.C. ALCARAZ¹ and R.Z. BARIEV^{1,2}

¹*Departamento de Física, Universidade Federal de São Carlos,
13565-905, São Carlos, SP Brazil*

²*The Kazan Physico-Technical Institute of the Russian Academy of Sciences,
Kazan 420029, Russia*

A new family of exactly solvable models is introduced. These models are generalizations of the XXZ chain where the distance among spins up (σ^z -basis) cannot be smaller or equal to t ($t = 0, 1, 2, \dots$). The case $t = 0$ recovers the standard XXZ chain. The coordinate Bethe ansatz is applied and the phase diagram is calculated. Exploring the finite-size consequences of conformal invariance the critical exponents are evaluated exactly at the critical regions of the phase diagram.

a

1 Introduction

The anisotropic $S = \frac{1}{2}$ Heisenberg model, or XXZ chain, is one of the most studied quantum spin system in statistical mechanics. Since its exact solution by Yang and Yang¹ this model has been considered to be the classic example of the success of the Bethe ansatz. In this paper, considering the XXZ chain as a prototype we introduce a new set of quantum chains which preserve the property of integrability.

The XXZ Hamiltonian on an L -site chain is given by

$$H(\Delta) = -\frac{1}{2} \sum_{i=1}^L (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+1}^z), \quad (1)$$

where $\sigma^x, \sigma^y, \sigma^z$ are spin- $\frac{1}{2}$ Pauli matrices, Δ is the anisotropy parameter and we assume periodic boundary conditions. If we identify, in the σ^z -basis, a spin up or spin down at a given site i as a particle or hole at this site, the XXZ Hamiltonian can be written equivalently as

$$H(\Delta) = - \sum_{i=1}^L P_0 (c_i^+ c_{i+1} + c_{i+1}^+ c_i + \Delta n_i n_{i+1}) P_0, \quad (2)$$

^aTo appear in the Proceedings of the Conference: Statistical Physics on the Eve of the Twenty-Fist Century

where c_i^+ and c_i are spinless creation and annihilation operators and $n_i = c_i^+ c_i$ is the density of particles at site i . The projector P_o in (1) forbids two particles to occupy the same position in space (fermion like). The models under consideration are obtained by a generalization of the operator P_0 and are given by

$$H(\Delta) = - \sum_{i=1}^L P_t (c_i^+ c_{i+1} + c_{i+1}^+ c_i + \Delta n_i n_{i+t+1}) P_t, \quad t = 0, 1, 2, \dots \quad (3)$$

where P_t now forbids two particles at distance less or equal to t ($0, 1, 2, \dots$). The static interaction, controlled by Δ , acts only when two particles are at the closest possible positions $(i, i+t+1)$. The case $t = 0$ recovers the standard XXZ chain (2) or (1). In terms of Pauli matrices we can also write the equivalent Hamiltonian

$$H_t(\Delta) = - \frac{1}{2} \sum_{i=1}^L P_t (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+t+1}^z) P_t, \quad (4)$$

where now P_t projects out from the Hilbert space the states (in the σ^z -basis) where two up spins ($\sigma_t = 1$) are at distance smaller or equal to t . In coming from (1) to (2) or from (3) to (4) we neglect a term proportional to a magnetic field. This term can be trivially added in the above Hamiltonian since the magnetization $\sum_{i=1}^L \sigma_i^z$ or the total number of particles $n = \sum_{i=1}^L n_i$ are good quantum numbers. In terms of Pauli matrices the operator $P_0 = 1$ and for $t > 0$ it is non-local and can be written in the σ^z -basis as

$$P_t = \prod_i \left[\frac{1}{2}(1 - \sigma_i^z) + \frac{1}{2}(1 + \sigma_i^z) \prod_{l=1}^t \frac{1}{2}(1 - \sigma_{i+l}^z) \right]. \quad (5)$$

The Hamiltonians (2), although looking artificial, may describe interesting physical situations where the states with up spins are associated with larger excluded volume than those of down spins. This is precisely the case when we consider the time fluctuations due to diffusion of molecules with size $t+1$ (in units of the lattice spacing parameter) on a lattice. Interpreting the master equation associated to this problem as a Schrödinger equation in Euclidean time,² the associated Hamiltonian is given by the isotropic ferromagnetic version of (3), namely $H_t(\Delta = +1)$. The anisotropic ferromagnetic version ($\Delta > 1$) is related to the asymmetric diffusion of the molecules of size $(t+1)$.³

2 The Bethe ansatz equations

It is important to observe that similarly to the XXZ chain the z -magnetization $m = \sum_i^L \sigma_i^z$, or the total number of up spins n (or the number of particles in (2) and (3)) is a conserved quantity for the Hamiltonian (4). Consequently the Hilbert space can be separated into block-disjoint sectors labelled by $n = 0, 1, \dots, L/(t+1)$. In a given sector n the wave functions of (4) can be written

$$\Psi = \sum_{x_1=1}^{\bar{L}} \dots \sum_{x_{i+1}=x_i+t+1}^{\bar{L}+i(t+1)} \dots \sum_{x_n=x_{n-1}+t+1}^L a(x_1, x_2, \dots, x_n) |x_1, x_2, \dots, x_n\rangle, \quad (6)$$

where $\bar{L} = L - n + 1 - t(n-1)$ and $a(x_1, \dots, x_n)$ are the amplitudes of the spin configurations $|x_1, x_2, \dots, x_n\rangle$, having spin down at the coordinates x_i ($i = 1, 2, \dots, n$). We can calculate the above amplitudes and the corresponding eigenenergy of (4) by applying the Bethe ansatz

$$a(x_1, x_2, \dots, x_n) = \sum_P A_{k_{p_1}, k_{p_2}, \dots, k_{p_n}} e^{i(k_{p_1}x_1 + \dots + k_{p_n}x_n)}, \quad (7)$$

where the summation is over all permutations $P = (p_1, p_2, \dots, p_n)$ and $A_{\{k\}}$ as well the wave numbers $\{k\} = k_1, k_2, \dots, k_n$ should be calculated from the eigenvalue equation for (4). The solution follows the standard coordinate Bethe ansatz.⁴ The validity of (7) for the case $n = 1$ is immediate due to the translational invariance of (4), and in this case the energies are $E_k = -2 \cos k - \frac{\Delta}{2}(L-4)$, with $k = 2\pi p/L$ ($p = 0, 1, 2, \dots, L-1$). The case of general n can be verified as in the standard XXZ chain ($t = 0$). The energies and momentum are given by

$$E_k = -2 \sum_{i=1}^n \cos 2k_i - \frac{\Delta}{2}(L-4n), \quad P = \sum_{i=1}^n k_i. \quad (8)$$

The wave-numbers $\{k_i\}$ are obtained by solving the Bethe ansatz equations

$$e^{ik_j L} = (-1)^{n-1} \prod_{l=1}^n e^{it(k_j - k_l)} \frac{1 - 2\Delta e^{ik_j} + e^{i(k_j + k_l)}}{1 - 2\Delta e^{ik_l} + e^{i(k_j + k_l)}}. \quad (9)$$

These equations at $t = 0$ give us the well-known Bethe ansatz equations for the XXZ chain. For $t > 0$ these equations in a given sector specified by n up spins and momentum $P = \frac{2\pi}{L}p$ ($p = 0, 1, \dots, L-1$) are the same as those of an XXZ chain also with n up spins but lattice size $L' = L - tn$, momentum $P' = \frac{2\pi}{L-tn}p \pmod{2\pi}$ and twisted boundary condition⁵

$$\sigma_{L+1}^x \pm i\sigma_{L+1}^y = e^{i\Phi}(\sigma_1^x \pm i\sigma_1^y), \quad \sigma_{L+1}^z = \sigma_1^z, \quad \Phi = \frac{2\pi}{L}p. \quad (10)$$

We will now study the phase diagram of these models. Calculating the eigen-spectra of (4) numerically by brute-force diagonalization on small lattices, we see that in absence of an external magnetic field the groundstate, for finite chains will belong to a sector where the magnetization $M = 2n - L$ depends on the lattice size L , exclusion parameter t and interaction Δ , as long as $t > 0$. This is distinct from the standard XXZ case ($t = 0$), where the groundstate is always in the half-filled sector $n = \frac{L}{2}$, with zero magnetization for all values of L and $\Delta < 1$. In order to have a well defined thermodynamic limit for given values of t and Δ it is necessary to work with fixed values of the magnetization per site $m = \frac{M}{L} = \frac{2n}{L} - 1$, or equivalently for fixed values of the density of up spins $\rho = \frac{n}{L}$. We can force the groundstate to belong to such a sector with $n = \rho L$ by adding in (4) an external magnetic field $-h \sum_{i=1}^L \sigma_i^z$, which will depend on the values of t , Δ and ρ .

It is important to remark that even when $\Delta = 0$ we do not have a completely free system for $t > 0$, since in (9) a phase connecting different values of k_j is always present. Nevertheless the Bethe-ansatz equations can be solved exactly for finite lattices, and is then instructive to calculate the eigenspectra at $\Delta = 0$.

3 The operator content at $\Delta = 0$

Let us consider $H_t(\Delta = 0, h)$, with a given value of the magnetic field $h = h(\rho)$ that produces a fixed density $\rho = \frac{n}{L}$ ($0 \leq \rho \leq \frac{1}{t+1}$) of up spins. Taking the logarithm of both sides of (9) we obtain

$$Lk_j = \pi(\rho L - 1) + \sum_{l=1}^{\rho L} t(k_j - k_l) + 2\pi m_j, \quad j = 1, 2, \dots, \quad (11)$$

where m_j ($j = 1, 2, \dots, n$) are integers. The solution of these equations are given by the set $\{k_j\}$,

$$k_j = \Theta + \frac{2\pi}{L(1 - t\rho)} m_j, \quad \Theta = \frac{\pi(\rho L - 1) - tP}{L(1 - \rho t)}, \quad j = 1, 2, \dots, n, \quad (12)$$

where $\{m_j\}$ are distinct integers and P is the momentum associated to the state

$$P = \sum_{j=1}^{\rho L} k_j = \frac{2\pi}{L} \tilde{p} \pmod{2\pi}, \quad \tilde{p} \in Z. \quad (13)$$

The groundstate has zero momentum and corresponds to the set $\{k_j\}$, where

$$k_j = \frac{2\pi}{L(1-t\rho)}j, \quad j = -\frac{\rho L-1}{2}, -\frac{\rho L-1}{2}+1, \dots, \frac{\rho L-1}{2}. \quad (14)$$

Using (14) in (8) we obtain the groundstate energy of (4), in the presence of an external magnetic field,

$$E_0(L, \rho) = -2 \sin\left(\frac{\pi\rho}{1-t\rho}\right) / \sin\left(\frac{\pi}{L(1-t\rho)}\right) - h(2\rho-1)L. \quad (15)$$

The magnetic field $h = h(\rho)$ is to be calculated by imposing that when $L \rightarrow \infty$, $e_\infty = \frac{E_0(L, \rho)}{L}$ will be a minimum as a function of ρ , i.e.

$$h = h(\rho) = \frac{t}{\pi} \sin\left(\frac{\pi\rho}{1-t\rho}\right) - \frac{1}{1-t\rho} \cos\left(\frac{\pi\rho}{1-t\rho}\right). \quad (16)$$

This result also tells us that in the absence of a magnetic field the groundstate should belong to the sector with magnetization $m_0 = \frac{M_0}{L} = 2\rho_0 - 1$, where $h(\rho_0) = 0$. In the case $t = 0$, $\rho_0 = \frac{1}{2}$ and $m_0 = 0$. For $t > 0$ we should solve $h(\rho_0) = 0$ numerically; for example, $\rho_0 = 0.3008443$, $\rho_0 = 0.2193388$ and $\rho_0 = 0.07354512$ for $t = 1, 2$ and 10 , respectively. As we can see these values cannot be predicted by symmetry arguments on (4) or (9). The leading finite-size corrections of (15) are given by

$$\frac{E^0(L, \rho)}{L} = e_\infty - \frac{\pi c}{6L^2} \xi + o(L^{-2}), \quad (17)$$

where

$$e_\infty = -2 \frac{1-\rho t}{\pi} \sin\left(\frac{\pi\rho}{1-\rho t}\right) - h(2\rho-1), \quad c\xi = \frac{2}{1-t\rho} \sin\left(\frac{\pi\rho}{1-t\rho}\right). \quad (18)$$

These relations are a clear indication that the model is critical for arbitrary values of t and ρ , since (17) is the expected behavior of a massless conformally invariant critical point with c and ξ being the conformal anomaly and sound velocity, respectively.⁶ The sound velocity can be calculated from the energy momentum dispersion relation. The lowest eigenenergy in the sector with $n = \rho L$ spins up, having momentum $\frac{2\pi}{L}\tilde{p}$ ($\tilde{p} = 1, 2, \dots$) can be obtained by choosing the particular solution of (11),

$$\begin{aligned} k_j &= \frac{2\pi}{L(1-t\rho)}j - \frac{2\pi t}{(1-t\rho)L^2}, \\ j &= -\frac{\rho L-1}{2}, -\frac{\rho L-1}{2}+1, \dots, \frac{\rho L-3}{2}, \frac{\rho L-1}{2} + \tilde{p}, \end{aligned} \quad (19)$$

and the corresponding energy is given by

$$E^{(\tilde{p})}(L, \rho) = E^0(L, \rho) + 4 \sin \left(\frac{\pi(\rho L + \tilde{p} - 1)}{(1 + t\rho)L} \right) \sin \left(\frac{\pi\tilde{p}}{L(1 - t\rho)} \right) + o(L^{-1}). \quad (20)$$

This gives us, for large L , a linear dispersion relation with sound velocity

$$\xi = \frac{2}{1 - t\rho} \sin \left(\frac{\pi\rho}{1 - t\rho} \right), \quad (21)$$

and consequently the conformal anomaly is $c = 1$ for all values of t and ρ . The conformal invariance of the infinite critical system, beyond relation (17), also predicts the finite-size corrections of excited states.⁷ For each primary operator, with dimension x_ϕ and spin s_ϕ , in the operator algebra of the critical system there exists an infinite tower of eigenstates of the Hamiltonian, whose energy $E_{m,m'}^\phi$ and momentum $P_{m,m'}^\phi$, in a periodic chain are given by

$$\begin{aligned} E_{m,m'}^\phi(L) &= E^0 + \frac{2\pi\xi}{L}(x_\phi + m + m') + o(L^{-1}), \\ P_{m,m'}^\phi &= (s_\phi + m - m')\frac{2\pi}{L}, \end{aligned} \quad (22)$$

where $m, m' = 0, 1, 2, \dots$. The groundstate corresponds to the identity operator $x_I = 0, s_I = 0$ and the tower of states $E_{m,m'}^{(\phi=I)} = \frac{2\pi}{L}(m - m')$ in our model are obtained by decreasing (increasing) the values of the quasi-momenta k_i in the set $\{k_i\}$ given in (14). After the calculation of the eigenenergies corresponding to many of these descendants we convince ourselves that the degeneracy of such levels are given by $d_{m,m'}^{\phi=I} = P(m)P(m')$ where $P(m)$ is the number of integer partitions of m . Another conformal dimension in the sector $n = \rho L$ can be obtained by taking negative (positive) values of k_j in the groundstate root configuration (14) and replacing them by positive (negative) values. For example, the lowest of these eigenenergies corresponds to the solution

$$\begin{aligned} k_j &= \frac{2\pi}{(1 - \rho t)L}(j - \rho t), \\ j &= -\frac{\rho L - 1}{2} + 1, -\frac{\rho L - 1}{2} + 2, \dots, \frac{\rho L - 1}{2} + 1, \end{aligned} \quad (23)$$

and has the value

$$E = -2 \cos \left(\frac{2\pi}{L} \right) \sin \left(\frac{\pi\rho}{1 - \pi\rho} \right) / \sin \left(\frac{\pi}{(1 - \rho t)L} \right) \quad (24)$$

and a macroscopic momentum $P = 2\pi\rho$. Comparing the finite-size corrections of (24) with (22) we see that this energy corresponds to an operator with dimension $x_{0,1}$ and spin $s_{0,1}$ given by

$$x_{0,\bar{m}} = (1 - \rho t)^2 \bar{m}^2, \quad s_{0,\bar{m}} = n\bar{m} = \rho L \bar{m}, \quad (25)$$

respectively. The dimensions $x_{0,\bar{m}}$, with $\bar{m}$ positive (negative) integer are obtained by taking $\bar{m}$ of the lowest (highest) values of k_j in (14) and replacing them by the lower (higher) available positive (negative) values. As in the case of the identity operator the descendants of the above operators, with dimension $x_{\bar{m}} + m + m'$ ($m, m' \in Z$) are obtained by increasing (decreasing) the positive (negative) values of $\{k_j\}$ in the configuration that produces $x_{0,\bar{m}}$. As before we convince ourselves that the degeneracy of these states is $P(m)P(m')$.

There exist also dimensions associated to eigenstates produced by the addition ($\bar{n} = -1, -2, \dots$) or subtraction ($\bar{n} = 1, 2, \dots$) of spins up in the ground-state sector. In this case $n = \rho L + \bar{n}$ and the lowest eigenenergy is obtained from the set $\{k_j\}$ with

$$k_j = \frac{2\pi}{L(1 - \rho t - \bar{n})} j, \quad j = -j_{\max}, -j_{\max} + 1, \dots, j_{\max} - 1, j_{\max}, \quad (26)$$

where $j_{\max} = \frac{\rho L + \bar{n} - 1}{2}$. These are zero-momentum eigenstates with energies

$$\begin{aligned} E^{\bar{n}} &= -2 \sin \left(\frac{\pi(\rho L + \bar{n})}{L(1 - \rho t) - \bar{n}} \right) / \sin \left(\frac{\pi}{L(1 - \rho t) - \bar{n}} \right) \\ &\quad - h(2(\rho L + \bar{n}) - L). \end{aligned} \quad (27)$$

The finite-size corrections, when compared with (22) give us the conformal dimensions

$$x_{\bar{n},0} = \frac{\bar{n}^2}{4(1 - t\rho)^2}. \quad (28)$$

In the general case, where we combine $\{k_j\}$ configurations which give (25) and (26) we obtain the conformal dimensions

$$x_{\bar{n},\bar{m}} = \bar{n}^2 X_p + \frac{\bar{m}^2}{4X_p}, \quad X_p = \frac{1}{4(1 - t\rho)^2} \quad \bar{n}, \bar{m} \in Z \quad (29)$$

of primary operators with spin $s_{\bar{n},\bar{m}} = (\rho L + \bar{n})\bar{m}$. If we calculate the eigenenergies which correspond to the descendants with dimensions $x_{\bar{n},\bar{m}}^{m,m'} = x_{\bar{n},\bar{m}} + m + m'$ we obtain their degeneracy $d_{\bar{n},\bar{m}}^{m,m'} = P(m)P(m')$. The dimensions

(29) are the same as the dimensions $x_{\bar{n},\bar{m}}^G = \frac{\bar{n}^2}{4\pi K} + \pi K \bar{m}^2$ of operators with vorticity $\bar{m}$ and spin-wave number $\bar{n}$ in a Gaussian model⁸ with coupling constant $K = (1 - \rho t)^2 / \pi$. The degeneracy $d_{\bar{n},\bar{m}}^{m,m'}$ imply that the operator content of $H_t(\Delta = 0)$ is given in terms of U(1) Kac-Moody characters. The situation is completely analogous with that of the standard XXZ chain ($t = 0$) in its massless regime.⁹ However, expression (29) shows different dependences on the density ρ (or magnetization), depending on the value of t . In particular for $\Delta = 0$ the dimensions are ρ -independent only in the free-fermion case $t = \Delta = 0$. The result (29) shows that the net effect of the parameter $t > 0$ at $\Delta = 0$ is similar to that of the anisotropy parameter $\Delta \neq 0$ in the standard XXZ chain ($t = 0$). In both cases the exponent X_p changes continuously with the density ρ of up spins. It is interesting to remark that when $\rho \rightarrow 0$ $X_p = 1/4$, for all values of t . This independence of t arises since the restrictions imposed by $t > 0$ are not important due to the small amount of up spins in this limit. The other limit of X_p at high density, $\rho \rightarrow 1/(t + 1)$, can also be understood as follows. If we start with the maximum density $\rho = 1/(t + 1)$ and we take out one spin up, the “hole” produced will have the same effective motion as that produced by adding $t + 1$ “particles” in the low-density limit. Then according to (28) we should obtain the limiting value $X_p = (t + 1)^2/4$.

4 The operator content at $\Delta \neq 0$

Let us consider the case where $\Delta \neq 0$. The Bethe ansatz equations (9) supplemented by numerical calculations give the same phase diagram for all values of t . For $\Delta > 1$ the groundstate is $(t + 2)$ -degenerate, where $(t + 1)$ states are in the sector with maximum number of up spins ($n = \frac{L}{1+t}$) and the last in the sector with no spins up ($n = 0$). For $\Delta < 1$ we expect a massless behavior similar to that in the XXZ chain. In fact, exploring the similarity of our equations for arbitrary t , with those of the XXZ chain ($t = 0$), we can apply the method of Ref. 10 to extract the finite-size corrections for arbitrary values of Δ and ρ . The dimensions we obtained are given by (29), where X_p is now given by

$$X_p = \frac{1}{4}(1 - t\rho)^{-2}\eta^{-2}(U_0), \quad (30)$$

where U_0 and $\eta(U)$ are obtained by solving some integral equations. These equations for $-1 \leq \Delta = -\cos \gamma \leq 1$ are given by

$$Q(U) = \frac{1}{2\pi} \frac{\sin \gamma}{\cos U - \cos \gamma} - \frac{1}{2\pi} \int_{-U_0}^{U_0} \frac{\sin(2\gamma)Q(U')}{\cosh(U - U') - \cos(2\gamma)} dU', \quad (31)$$

$$1 = \eta(U) + \frac{1}{2\pi} \int_{-U_0}^{U_0} \frac{\sin(2\gamma)\eta(U')}{\cosh(U - U') - \cos(2\gamma)} dU', \quad (32)$$

while for $\Delta = -\cosh \gamma < -1$ they are given by

$$Q(U) = \frac{1}{2\pi} \frac{\sinh \lambda}{\cosh \lambda - \cos U} - \frac{1}{2\pi} \int_{-U_0}^{U_0} \frac{\sinh(2\lambda) Q(U')}{\cosh(2\lambda) - \cos(U - U')} dU', \quad (33)$$

$$1 = \eta(U) + \frac{1}{2\pi} \int_{-U_0}^{U_0} \frac{\sinh(2\lambda) \eta(U')}{\cosh(2\lambda) - \cos(U - U')} dU'. \quad (34)$$

In Eqs. (31)-(34) U_0 is calculated by imposing

$$\int_{-U_0}^{U_0} Q(U) = \begin{cases} \frac{\rho}{1-t\rho}, & 0 \leq \rho \leq \frac{1}{2+\rho}, \\ \frac{1-(t+1)\rho}{1-t\rho}, & \frac{1}{2+\rho} \leq \rho \leq \frac{1}{1+\rho}. \end{cases} \quad (35)$$

The value of X_p is obtained by numerically solving Eqs. (31)-(35). In Fig. 1 we plot X_p at $\Delta = -\frac{\sqrt{2}}{2}$ for $t = 0, 1, 2$ and 3, while in Fig. 2 X_p is shown for $t = 2$ and some values of Δ ($1 \leq \Delta \leq 1$).




Figure 1: The value of X_p as a function of $\rho = \frac{n}{L}$ at $\Delta = -\frac{\sqrt{2}}{2}$, for $t = 0$ (a), $t = 1$ (b), $t = 2$ (c) and $t = 3$ (d).

These results coincide with those calculated earlier for the standard XXZ chain ($t = 0$).¹⁰ At low density ($\rho \rightarrow 0$) and high density ($\rho(t+1) \rightarrow 1$) the exponent $X_p = 1/4$ and $X_p = (t+1)^2/4$, respectively. These results are the same as those of the $\Delta = 0$ case, since in these limits the distribution of “particles” at $\rho \rightarrow 0$ or “holes” at $\rho \rightarrow 1/(t+1)$ are so diluted as to render the effect of the interaction term Δ negligible. From (30)-(34) it is possible to relate $X_p(t, \Delta)$ for different values of t and ρ . In particular at the point $\rho = 1/(2+t)$, which corresponds to the “half-filled” density for these models,




Figure 2: The value of X_p as a function of $\rho = \frac{n}{t}$ for $t = 2$ and anisotropy $-1 \leq \Delta = -\cos \gamma \leq 1$, with (a) $\gamma = \frac{\pi}{10}$, (b) $\gamma = \frac{\pi}{4}$, (c) $\gamma = \frac{\pi}{3}$, (d) $\gamma = \frac{\pi}{2}$ and (e) $\gamma = \frac{2\pi}{3}$.

we can show that

$$x_p(t, \frac{\rho}{2+t}) = \frac{1}{4}(t+2)^2 X_p(t=0, \rho = \frac{1}{2}) = \frac{1}{8}(t+2)^2(\pi - \gamma), \quad (36)$$

for $-1 \leq \Delta = -\cos \gamma \leq 1$, where we have used the known results⁵ for the XXZ chain in the absence of an external magnetic field.

In Fig. 3 we plot X_p for $t = 3$ and for some values of $\Delta \leq -1$, while in Fig. 4 X_p is shown at $\Delta = -2$ for $t = 0, 1, 2$ and 3.

As before in the low density ($\rho \rightarrow 0$) and high density ($\rho \rightarrow 1/(t+1)$) limits we recover the $\Delta = 0$ case. In Fig. 3 we also notice that at the “half-filling” density $\rho = 1/(2+t)$ the exponent $X_p = (t+2)^2/4$, independently of the value of Δ . A physical explanation for this behavior is the following. Strictly speaking at the “half-filling” point $\rho = 1/(t+2)$ the model has a gap for $\Delta < -1$, as long as the magnetic field is zero. When the magnetic field increases, reaching a critical value $h = h_c(\Delta, t)$ this gap vanishes. Therefore the main physical effect at $\rho = 1/(2+t)$ is just the vanishing of the gap and not the interaction Δ itself. In the limit $\Delta \rightarrow -\infty$ and $\rho < 1/(2+t)$ the model with a given value of t behaves effectively as a model with $t' = t+1$ and $\Delta = 0$, since the spins up have infinite repulsion and will be located in positions with no interactions among them (distances greater than $t+1$). This can also be verified directly by setting $\Delta \rightarrow -\infty$ in the Bethe ansatz equations (9). Consequently for $\Delta \rightarrow -\infty$ and $\rho < 1/(2+t)$ we should have $X_p = 1/[4(1-\rho(t+1))^2]$. In particular at the “half-filling” density $\rho = 1/(2+t)$ we obtain the value $X_p = (t+2)^2/4$. In Fig. 3 the limiting curve ($\Delta \rightarrow -\infty$),




Figure 3: The curves $X_p(\rho)$ for $t = 3$ and (a) $\Delta = -20$, (b) $\Delta = -4$, (c) $\Delta = -2$. The expected curve at $\Delta \rightarrow -\infty$ $X_p = 1/[4(1 - 4\rho)^2]$ is also shown.




Figure 4: The curves $X_p(\rho)$ at $\Delta = -2$ for (a) $t = 0$, (b) $t = 1$, (c) $t = 2$ and (d) $t = 3$.

$X_p = 1/4(1-4\rho)^2$, for $t = 3$, can be compared with that of $\Delta = -20$, obtained by solving (30) and (33)-(35). For $\Delta \rightarrow -\infty$, but $\rho > 1/(2+t)$ this argument does not apply and we should recover the value $X_p = (t+1)^2/4$ at $\rho = 1/(t+1)$. We believe the results reported here in the $\Delta < -1$ region are unknown even for the standard XXZ chain ($t = 0$).

5 Generalizations

Before concluding let us discuss an extension of our results. We only considered the cases where $t \geq 0$ but what should be the quantum chain that would correspond to $t = -1$? In terms of particles, like in (2) this corresponds to the situation where we can have an arbitrary number of particles on a site. In fact we were able to find an exactly integrable quantum chain in this class, namely

$$\begin{aligned} H_{-1}(\Delta) = & - \sum_{j=1}^L \left\{ \sum_{n=0}^{\infty} \sum_{m=n+1}^{\infty} (E_j^{n,m} E_{j+1}^{m,n} + E_j^{m,n} E_{j+1}^{n,m}) \right. \\ & \left. + \sum_{n=0}^{\infty} \Delta(n-1) E_j^n E_{j+1}^n \right\}, \end{aligned} \quad (37)$$

where $E^{n,m}$ are ∞ -dimensional matrices located at the lattice sites with elements $E^{n,m}_{i,j} = \delta_{n,i} \delta_{m,j}$. The Bethe ansatz equations are given by (9) with $t = -1$. This Hamiltonian commutes with the operator

$$\hat{N} = \sum_{j=1}^L \sum_{n=1}^{\infty} n E_j^{n,n}, \quad (38)$$

which gives the total number of particles. The Hamiltonian (37) in a given sector with N particles can be written in terms of spin- S $SU(2)$ matrices with $S = (N-1)/2$.

Acknowledgments

This work was supported in part by CNPq and FINEP – Brazil, and also by the Russian Foundation of Fundamental Investigations under Grant No.RFFI 97-02-16146.

References

1. C.N. Yang and C.P. Yang, *Phys.Rev.* **150**, 320, 327 (1966).

2. See, e.g, F.C. Alcaraz, M. Droz, M.Henkel and V. Rittenberg, *Ann. Phys. (NY)* **230**, 250 (1994).
3. F.C. Alcaraz and R.Z. Bariev, to be published.
4. See, e.g., R.J. Baxter, *Exactly Solved Models in Statistical Mechanics* (Academic Press, New York, 1982).
5. F.C. Alcaraz, M.N. Barber and M.T. Batchelor, *Ann. Phys. (NY)* **182**, 280 (1988); *Phys. Rev. Lett.* **58**, 771 (1987).
6. H.W.J. Blöte, J.L. Cardy and M.P. Nightingale, *Phys. Rev. Lett.* **56**, 742 (1986); I. Affleck, *Phys. Rev. Lett.* **56**, 746 (1986).
7. J.L. Cardy, in *Phase Transitions and Critical Phenomena* Vol.11, C. Domb and J.L. Lebowitz eds. (Academic Press, London, 1987).
8. L. Kadanoff and A. Brown, *Ann. Phys. (NY)* **121**, 318 (1979).
9. F.C. Alcaraz, M. Baake, U. Grimm and V. Rittenberg, *J. Phys. A* **21**, L117 (1988).
10. F. Woynarovich, H.P. Eckle and T.T. Truong, *J. Phys. A* **22**, 4027 (1989).